

A new male-killing parasitism: *Spiroplasma* bacteria infect the ladybird beetle *Anisosticta novemdecimpunctata* (Coleoptera: Coccinellidae)

M. C. TINSLEY* and M. E. N. MAJERUS

Department of Genetics, University of Cambridge, Downing Street, Cambridge CB2 3EH, UK

(Received 3 October 2005; revised 28 November 2005; accepted 1 December 2005; first published online 3 February 2006)

SUMMARY

Whilst most animals invest equally in males and females when they reproduce, a variety of vertically transmitted parasites has evolved the ability to distort the offspring sex ratios of their hosts. One such group of parasites are male-killing bacteria. Here we report the discovery of females of the ladybird *Anisosticta novemdecimpunctata* that produced highly female-biased offspring sex ratios associated with a 50% reduction in egg hatch rate. This trait was maternally transmitted with high efficiency, was antibiotic sensitive and was infectious following experimental haemolymph injection. We identified the cause as a male-killing *Spiroplasma* bacterium and phylogenetic analysis of rDNA revealed that it belongs to the *Spiroplasma ixodetis* clade in which other sex ratio distorters lie. We tested the potential for interspecific horizontal transfer by injection from an infected *A. novemdecimpunctata* line into uninfected individuals of the two-spot ladybird *Adalia bipunctata*. In this novel host, the bacterium was able to establish infection, transmit vertically and kill male embryos.

Key words: male-killing, Coccinellidae, *Spiroplasma*, *Anisosticta novemdecimpunctata*, *Adalia bipunctata*, horizontal transfer.

INTRODUCTION

Endosymbiont microbes that disrupt host reproductive biology are common parasites of arthropods (Bandi *et al.* 2001). These reproductive parasites obligately inhabit host tissues and achieve transmission only within the egg when female hosts reproduce. To such vertically transmitted bacteria, males are therefore an evolutionary dead end (Hurst, 1991). Transmission to offspring is typically not perfect and many infections impose fitness costs on female hosts. Nevertheless, reproductive parasites can proliferate through populations by biasing host offspring sex ratios in favour of the transmitting sex (females). Sex ratio distortion may be achieved by 3 different mechanisms: male-killing, feminization, and parthenogenesis induction. An alternative reproductive manipulation strategy to sex ratio distortion is cytoplasmic incompatibility (Bandi *et al.* 2001).

Male-killing bacteria cause male offspring death during embryonic development, resulting in low egg hatch rates in clutches laid by infected females. This is probably the most common mechanism by which reproductive parasites achieve sex ratio distortion and is known only from insects (Hurst and Jiggins, 2000). The phenotype is selected for when the death

of male embryos elevates the immediate fitness of sibling females which also harbour the bacterium (Hurst, 1991). This occurs if male death reduces the opportunities for deleterious sibling inbreeding, reduces antagonistic sibling interactions (e.g. egg cannibalism, competition for resources), or if neonate female survival is enhanced by the consumption of dead male embryos within the brood (Hurst and Majerus, 1993). The spread of male-killers in host populations is therefore crucially dependent on host ecology, in addition to genetic interactions between insect and parasite.

The bacterial genus *Wolbachia* has been considered exceptional amongst reproductive parasite taxa. Infections are very common; most estimates suggest that 17–30% of arthropod species are hosts (Werren, Windsor and Guo, 1995; Werren and Windsor, 2000). Initial reports indicated that *Wolbachia* was unique in its ability to perform all 4 types of parasitic reproductive manipulation. However, more recently this has been challenged by the findings that members of the Cytophaga-Flavobacterium-Bacteriodes phylum can also induce all 4 phenotypes (Hurst *et al.* 1997; Weeks, Marec and Breeuwer, 2001; Zchori-Fein *et al.* 2001; Hunter, Perlman and Kelly, 2003), although the proportion of species infected appears lower than for *Wolbachia* (Zchori-Fein and Perlman, 2004). In contrast to the limited taxonomic range of bacteria known to have adopted parthenogenesis induction, feminization and cytoplasmic incompatibility, a

* Current address for correspondence: School of Biological and Environmental Sciences, University of Stirling, Stirling FK9 4LA, UK. Tel: 00 44 (0) 1786 467840. Fax: 00 44 (0) 1786 467843. E-mail: matthew.tinsley@stir.ac.uk

phylogenetically diverse array of microbes is responsible for male-killing. Male-killers have been reported from 5 different bacterial taxa: *Spiroplasma*, γ -proteobacteria, *Rickettsia*, Flavobacteria and *Wolbachia* (Hurst and Jiggins, 2000).

Majerus and Hurst (1997) suggested that the aphidophagous Coccinellidae are particularly prone to invasion by male-killers due to their ecology: indeed, over 50% of species are estimated to be infected (Hurst, Jiggins and Majerus, 2003). Thus coccinellids are ideal organisms for investigating the diversity of symbionts responsible for male-killing. The majority of known host species are generalists, with broad ecological requirements. In contrast, *Anisosticta novemdecimpunctata* is a habitat specialist found only on emergent aquatic plants or terrestrial vegetation immediately adjacent to water bodies (Majerus, 1991). Majerus and Hurst (1997) outlined four criteria which may predispose a coccinellid species to male-killing infection: it should be aphidophagous, lay eggs in clutches, indulge in larval sibling egg cannibalism, and starvation should be a major cause of neonate larval mortality. *Anisosticta novemdecimpunctata* fulfils the first three of these, and whilst the latter has not been investigated, it seems probable it applies. Hence, we predicted that *A. novemdecimpunctata* would be an ecologically suitable host for male-killer infection. Our discovery of females that produced only daughters provided an opportunity to investigate the nature and cause of this sex ratio trait.

MATERIALS AND METHODS

The sex ratio trait

Anisosticta novemdecimpunctata were collected from *Phragmites* and *Typha* dominated aquatic habitats. Two English locations were sampled, one in the East (Cambridge: grid reference, TL 434587) and one in the Southwest (Chew Lake, Somerset: grid reference ST 575592) in October 2001 and March 2002 respectively. Adult beetles were maintained in the laboratory in 9 cm diameter Petri dishes at 21 °C under a 24 h L photo-period and fed pea aphids (*Acyrtosiphon pisum*) and artificial diet (Majerus *et al.* 1989). Petri dishes were changed daily; those with egg clutches were retained, eggs counted and hatch rates of clutches subsequently determined. Larvae were reared and sexed when adult under carbon dioxide anaesthesia. Male and female *A. novemdecimpunctata* are distinguished by a prominent anteriorly-directed groove in the posterior-most ventral abdominal segment of females and a clear ingression of the same region in males (Randall, Majerus and Forge, 1992). Female-biased broods were identified and inheritance investigated in the F₁ generation. To determine whether sex ratio distortion was bacterial in origin, F₁ females were treated

with the antibiotic tetracycline hydrochloride in golden syrup (10% w/w) for 4 h daily over 10 days (Hurst, Majerus and Walker, 1992). Eggs were collected before, during and after this treatment and resulting larvae reared to adulthood. Golden syrup alone does not influence offspring sex ratios in other coccinellids (Hurst *et al.* 1992). To confirm this, 1 F₁ female was fed only golden syrup as a control. Offspring samples produced during treatment were preserved for molecular analysis.

Infectivity of the sex ratio trait

Donor *A. novemdecimpunctata* from all-female lineages were fed excess aphids for 1 week to promote ovary development and ensure adequate tissue volume. *Adalia bipunctata* from equitable sex ratio lines and uninfected by male-killers were reared to pupal stage to act as recipients. *Adalia bipunctata* was employed because its larger size facilitated injection and laboratory rearing is more successful than for *A. novemdecimpunctata*. Following the method of Hurst *et al.* (1999b), donor *A. novemdecimpunctata* abdomens were removed and homogenized in sterile 0.7% NaCl solution (10 μ l per abdomen) on ice. Injection needles were pulled from glass capillaries (1.1 mm internal diameter) and attached to a 10 μ l Hamilton syringe. Recipient *A. bipunctata* pupae were injected with 1 μ l of homogenate adjacent to the dorsal midline, between abdominal segments 2 and 3. Following pupal eclosion, males were discarded and females maintained for 1 month to permit bacterial replication, then mated and bred.

Identification of the bacterium responsible

Following the method of Jiggins *et al.* (2000), a modified version of the Chelex 100 protocol was employed to extract genomic DNA. Beetle abdomens were homogenized in buffer containing 200 μ l of sterile distilled water, 0.01 g Chelex-100 ion-exchange resin (Bio-Rad), 7 μ l of 1 M dithiothreitol and 1 μ l of proteinase K (20 mg ml⁻¹), then digested for 70 min at 56 °C. Extractions were subsequently incubated at 96 °C for 10 min and centrifuged at 16 000 g for 5 min. Bacterial presence was investigated by diagnostic PCR using primers specific for male-killing bacterial taxa: *Rickettsia* RSSUF–RSSUR (Schulenburg *et al.* 2001), *Wolbachia* wsp81f–wsp691r (Zhou, Rousset and O'Neill, 1998), Flavobacteria FL1–FL2 (Hurst *et al.* 1997) and Mollicutes MGSO–Ha–In-1 (van Kuppeveld *et al.* 1992; Hurst *et al.* 1999b). Mitochondrial DNA primers J2630–N3014 (Schulenburg *et al.* 2002) were used to verify the presence of amplifiable template and samples not yielding product were discarded. PCR reactions were run using a premix containing 12.6 μ l of sterile distilled water, 2 μ l of 10 \times PCR reaction buffer (Sigma), 2 μ l of MgCl₂ (25 mM; Sigma),

2 μ l of dNTPs (2 mM per nucleotide: Bioline), 0.2 μ l of each primer (50 pM) and 0.6 μ l of REDTaqTM DNA polymerase (1 U μ l⁻¹: Sigma). DNA extraction supernatant (0.5 μ l) was added and reactions run in either a Techne (Progene and Genius models) or Hybaid (OmniGene) thermal cycler including appropriate positive and negative controls. For primers wsp81f-wsp691r, FL1-FL2 and J2630-N3014, cycle conditions were: 2 min at 95 °C, then 35 cycles of 20 sec at 95 °C, 1 min at 60 °C and 1 min at 72 °C, followed by 10 min at 72 °C. For primers RSSUF-RSSUR and MGSO-Ha-In-1, cycle conditions were: 2 min at 95 °C, then 35 cycles of 20 sec at 95 °C, 30 sec at 57 °C and 30 sec at 72 °C, followed by 10 min at 72 °C. PCR products were electrophoresed on 1% agarose gels containing 0.5 ng ml⁻¹ ethidium bromide in 1× TAE buffer, then visualized and photographed under UV light.

The bacterial 16S rDNA gene and adjoining internally transcribed spacer (ITS) region were amplified from 2 all-female lineages in 2 overlapping fragments using primer pairs 27f-MGSO and Ha-In-1-SPITS-N2, specific for the Mollicutes and *Spiroplasma* genus respectively (van Kuppeveld *et al.* 1992; Hurst *et al.* 1999b). Both strands were sequenced using the PCR primers and one internal primer (AnSpInt1r: 5'-GTT TGG GCC GTG TCT CAT C-3'). Products were electrophoresed, bands excised and DNA purified using GenEluteTM gel extraction columns (Sigma) following manufacturer's instructions. Sequencing employed the ABI PRISM Dye Terminator cycle sequencing ready reaction kit (Perkin Elmer) and an ABI 384 automated sequencer (Perkin Elmer). Sequence homology was first investigated by BLAST search. The *A. novemdecimpunctata* sequence (EMBL Accession number AM087471) was then manually aligned with published *Spiroplasma* ribosomal sequences: 3 male-killing and 3 non-male-killing spiroplasmas from the *Spiroplasma ixodetis* clade and 2 sequences from the *Spiroplasma citri* group. The variable ITS region was unalignable for more distantly related sequences, therefore 2 data sets were separately investigated using PAUP* version 4.0b8 (Swofford, 1998). Initially, only the 16S rDNA region of all sequences was studied, followed by the full-length sequence for members of the *S. ixodetis* clade to which this novel *Spiroplasma* belongs. Tree topology was evaluated by maximum likelihood using a heuristic search, branch swapping by nearest neighbour interchange and employing the HKY85+G substitution model (Hasegawa, Kishino and Yano, 1985). Model parameter values were calculated from an initial tree constructed using maximum parsimony and a tree bisection and reconnection branch swapping protocol. Statistical support for trees was sought by performing 1000 nonparametric bootstrap replicates. MEGA version 3.1 (Kumar, Tamura and Nei, 2004) was employed for sequence comparisons, calculating

Table 1. Breeding data for two UK samples of *Anisostica novemdecimpunctata* from Cambridge (Wa/) and Chew (Chew/)

(Lines displaying male-killing marked in bold. Egg hatch rates (proportion hatched) and offspring sex ratios (proportion male) are given (sample sizes in parentheses). Asterisks denote significant female-biased deviations from equitable sex ratios (Fisher's exact test, Bonferroni corrected for 26 multiple comparisons). Results of diagnostic PCR tests using Mollicute specific primers are indicated.)

Line number	Egg hatch rate	Offspring sex ratio	Mollicute specific PCR
Wa/2	0.28 (390)	0.00 (48)	+ve *
Wa/3	0.66 (235)	0.36 (25)	-ve
Wa/4	0.61 (165)	0.46 (13)	-ve
Wa/5	0.59 (62)	0.63 (8)	-ve
Wa/6	0.30 (103)	0.00 (27)	+ve *
Wa/7	0.13 (280)	0.00 (27)	+ve *
Wa/12	0.37 (140)	0.40 (55)	-ve
Wa/13	0.13 (65)	0.00 (3)	+ve
Wa/14	0.38 (115)	0.61 (54)	-ve
Wa/15	0.32 (98)	0.51 (43)	-ve
Wa/16	0.37 (43)	0.73 (11)	-ve
Wa/17	0.37 (123)	0.61 (72)	-ve
Wa/18	0.40 (203)	0.46 (100)	-ve
Chew/1	0.41 (145)	0.00 (52)	+ve *
Chew/2	0.64 (136)	0.36 (78)	-ve
Chew/3	0.60 (212)	0.53 (118)	-ve
Chew/4	0.57 (126)	0.51 (61)	-ve
Chew/5	0.26 (140)	0.00 (31)	+ve *
Chew/6	0.68 (159)	0.61 (89)	-ve
Chew/7	0.62 (175)	0.49 (101)	-ve
Chew/8	0.42 (218)	0.51 (63)	-ve
Chew/9	0.25 (145)	0.00 (40)	+ve *
Chew/10	0.32 (147)	0.51 (67)	-ve
Chew/11	0.36 (150)	0.45 (74)	-ve
Chew/12	0.46 (158)	0.48 (65)	-ve
Chew/13	0.31 (201)	0.04 (68)	+ve *

number of nucleotide changes and Kimura 2-parameter distances (using pair-wise gap deletion).

RESULTS

Sex ratio trait

Offspring were reared from 26 females, 13 from each site; mean offspring number per female was 53.6 (range 3–118). Progeny sex ratios of 7 females differed significantly from equity (Fisher's exact tests, Bonferroni corrected for multiple comparisons) (Table 1). Sex ratio biases were always accompanied by low egg hatch rates, indicative of male-killing (Table 1). A further female (Wa/13) exhibiting a low egg hatch rate produced only 3 offspring before dying, all were female. These 8 females had 293 female and 3 male offspring in total. Their mean egg hatch rate was 0.24 ($n=8$), exactly half that of the remaining females (0.48, $n=18$): hatch rates of these groups differed significantly (Mann Whitney U Test, $U_{(n=8,18)}=136$; $P=0.0004$).




Fig. 1. Offspring sex-ratios produced by *Spiroplasma*-infected F₁ females of *Anisostica novemdecimpunctata*. Vertical lines delimit an antibiotic treatment period during which females 2.2, 6.1, 6.2 and 6.3 received tetracycline in golden syrup (treat.), female 6.4 was a control and received only golden syrup (cont.). Each data-point represents mean offspring production over a 4-day period, numbers (*n*) above each figure provide sample sizes. Females 6.1 and 6.2 died before the experiment finished. Data for each female before and after treatment were pooled and sex ratios compared to those produced during treatment. *P*-values represent significance of sex ratio increases during treatment (Fisher's exact test, Bonferroni corrected for 5 comparisons).

DNA was extracted from 1 F₁ female from 2 biased (Wa/6 and Wa/2) and 2 equitable ratio lines (Wa/3 and Wa/4) and PCR tests performed. None produced amplification products using *Rickettsia*, *Wolbachia* or *Flavobacteria* specific primers. Both sex ratio lines did produce a product of approximately 430 bp with Mollicute-specific primers MGSO-Ha-In-1, but the 2 unbiased lines did not. The association between this bacterium and the sex ratio trait was assessed across all lines. PCR tests on either the parental female or 6 F₁ females revealed a perfect association

between this Mollicute bacterium and sex ratio distortion (Table 1). The single male offspring from sex ratio line Chew/13 that was tested carried the infection. Both population samples had a bacterial prevalence of 31% (*n* = 13; 95% CI = 6.6–52.0%).

Trait inheritance and antibiotic sensitivity

The inheritance of the sex ratio trait was investigated in 2 infected lines (Wa/2 and Wa/6). Eggs were collected from 5 F₁ females for approximately 25 days,




Fig. 2. Hatch rates of eggs produced by *Spiroplasma*-infected F₁ females of *Anisostica novemdecimpunctata*. Vertical lines delimit an antibiotic treatment period during which females 2.2, 6.1, 6.2 and 6.3 received tetracycline in golden syrup (treat.), female 6.4 was a control and received only golden syrup (cont.). Each data-point represents mean egg production over a 4-day period, numbers (*n*) above each figure provide sample sizes. Females 6.1 and 6.2 died before the experiment finished. Data for each female before and after treatment were pooled and hatch rates compared to those produced during treatment. *P*-values represent significance of hatch rate increases during treatment (Fisher's exact test, Bonferroni corrected for 5 comparisons).

before 4 were treated with antibiotic. Prior to treatment, all individuals produced only female offspring (mean = 30.4, range 16–50), accompanied by low hatch-rates (mean = 0.26, range 0.18–0.34). During and immediately following treatment, 3 of 4 females produced male offspring (range 1–9) (Fig. 1). Also, notably elevated hatch-rates occurred in 3 cases (Fig. 2). Sex ratio and hatch rate increases during antibiotic treatment were significant in 2 of 4 and 3 of 4 cases respectively (Figs 1 and 2). Alleviation of

male-killing symptoms was temporary: complete male mortality returned when antibiotic treatment ceased. The control line that received only golden syrup displayed low hatch-rates and produced solely female offspring throughout. All offspring (F₂) produced prior to antibiotic treatment that were tested were infected (*n* = 15) as were 22 F₂ females and 11 F₂ males derived afterwards. All control line offspring tested (6 before and 6 after treatment) were also infected.

Table 2. Breeding data for six *Adalia bipunctata* females injected with abdominal homogenate from female-biased *Anisostica novemdecimpunctata* lineages (sample sizes in parentheses)

(All *A. bipunctata* offspring sex ratios deviate significantly from equity (Fisher's exact test, Bonferroni corrected for 6 comparisons). Male-killing occurred in the novel species but was not complete.)

Line number	Egg hatch rate	Offspring		Offspring sex ratio
		Female	Male	
2sp/An/2	0.51 (280)	98	5	0.05
2sp/An/3	0.45 (249)	101	0	0.00
2sp/An/4	0.66 (247)	83	34	0.29
2sp/An/5	0.53 (530)	161	2	0.01
2sp/An/6	0.23 (151)	28	0	0.00
2sp/An/7	0.68 (269)	100	7	0.07

Trait infectivity

Abdomen homogenate of infected *A. novemdecimpunctata* females was injected into *A. bipunctata* pupae derived from 2 females producing equitable offspring sex ratios (0.46, $n=69$; 0.45, $n=83$) and carrying no known sex ratio distorting bacteria. Six females survived and laid eggs: all produced significantly female-biased offspring sex ratios (Fisher's exact tests, Bonferroni corrected for multiple comparisons) (Table 2). However, only 2 had exclusively female progeny and 1 produced numerous males (34 of 117 offspring). The mean egg hatch rate from these 6 injected *A. bipunctata* females was 0.51 (range 0.23–0.68) (Table 2). All male and female offspring tested were infected by the Mollicute bacterium ($n=34$ males and 9 females, representing all 6 lines).

Phylogenetic affiliation

The 16S and ITS rDNA region was sequenced for the *A. novemdecimpunctata* male-killer from 2 females providing 1672 bases for analysis and achieving approximately 80% double-stranded coverage. One site producing double peaks in the sequence of both forward and reverse strands for both individuals was ignored. The *A. novemdecimpunctata* Mollicute belonged to the *Spiroplasma* genus. Tree construction using maximum likelihood placed the male-killer in the *Spiroplasma ixodetis* clade, which is strongly evolutionarily divergent from the *Spiroplasma citri* clade containing the *Drosophila* male-killing spiroplasmas (SROs). Nonparametric bootstrapping provided no support for any branches except the *S. ixodetis* – *S. citri* clade divergence. All *S. ixodetis* clade sequences had very close homology (less than 1.5% difference), but were considerably divergent (more than 15%) from those in the *S. citri* clade (Table 3).

DISCUSSION

Male-killing bacteria are common parasites of insects where they distort population sex ratios in favour of females (Hurst *et al.* 2003). Here we report a new male-killing symbiosis; between a coccinellid beetle and a *Spiroplasma* bacterium. A number of other coccinellids are hosts to male-killers (Hurst and Jiggins, 2000), thus our discovery lends support to the suggestion of Majerus and Hurst (1997) that coccinellid ecology predisposes this group to invasion by male-killing parasites. The identification of a *Spiroplasma* as the bacterium responsible brings the number of coccinellid male-killers known of this bacterial genus to 3 (Hurst *et al.* 1999b; Majerus *et al.* 1999).

The male-killing trait was maternally inherited in all cases tested and transmission demonstrated across 3 host generations. In other coccinellids male-killer transmission occurs transovarially (Hurst *et al.* 1999a). The possibility of paternal transmission was not investigated here due to mortality of the 3 male offspring surviving from sex ratio lines, but seems unlikely. One F₁ male that was tested carried the infection, therefore its survival was not due to complete transmission failure. It seems probable that surviving males received low bacterial densities, permitting embryonic survival. *Spiroplasma* transmission was perfect, or near-perfect, under laboratory conditions, as observed for other coccinellid male-killers of this genus (Majerus *et al.* 1998; Hurst *et al.* 1999b). The apparent lack of inefficient transmission questions the mechanism for maintenance of this infection at moderate (31%) prevalence in the populations studied. In the majority of male-killing symbioses uninfected females are constantly generated by bacterial transmission failure, preventing parasite fixation and host extinction (Hurst and Jiggins, 2000). It is possible that field environmental factors may reduce transmission rates in natural populations.

The partial alleviation of male-killing symptoms during antibiotic treatment provides compelling evidence for the bacterial origin of this trait. Complete concordance of the sex ratio condition with the presence of the *Spiroplasma* supports this. The absence of a complete cure during antibiotic treatment was probably due to insufficient treatment length or dose. However, tetracycline exposure caused ladybird deterioration and 2 died, administering increased doses might have caused further mortality. Another *Spiroplasma* male-killer similarly proved difficult to cure by antibiotics (Majerus *et al.* 1998). Survival of infected male offspring following treatment suggests tetracycline either reduced bacterial density below the threshold required for embryo death, or disrupted male-killing directly. Dyer, Minhas and Jaenike (2005) found maternal bacterial density was inversely correlated with male

homology existed between *S. ixodetis* and this new *Spiroplasma*; furthermore the increasing numbers of bacteria allied to this clade provide a considerable number of comparators.

Anisosticta novemdecimpunctata differs from most previously described coccinellid male-killer hosts by being highly habitat specific (Majerus, 1991). Many aphidophagous coccinellid larvae starve shortly after hatching because they fail to find aphid prey (Hurst and Majerus, 1993). Killing males benefits the bacterium partly because infected female larvae consume dead embryos in their clutch, extending aphid search opportunities (Hurst and Majerus, 1993). In generalist coccinellid species adult females migrate between plant taxa during the reproductive season, ovipositing near productive aphid colonies where larval survival is relatively high (Sloggett and Majerus, 2000). Thus bacterial fitness returns of larval egg consumption may be small. However, habitat specialists are restricted to a narrow range of plants, more frequently laying eggs where aphids are scarce and bacterial fitness high. Therefore we predict male-killer prevalence may be greater in specialist than generalist hosts. Nonetheless, prevalence in both *A. novemdecimpunctata* populations was 31% and within the range typical for coccinellid male-killers (Hurst and Jiggins, 2000). Comparisons across more populations and species will be necessary to further test this hypothesis.

This study has revealed the existence of a new male-killing *Spiroplasma* parasite, closely phylogenetically allied to the other non-dipteran male-killers of this genus. Hurst *et al.* (2003) reviewed male-killer diversity across all insect species and concluded that the currently diminishing rate of discovery of new male-killing bacterial clades perhaps suggests that the majority have now been identified. That this male-killer also belongs to the *Spiroplasma* genus supports their argument. The present findings provide further demonstration of the widespread occurrence of male-killing as a strategy employed by a variety of parasitic agents in insect hosts.

We thank Frank Jiggins for much helpful advice and assistance and are grateful to Ian Wright and Dennis Farrington for technical support. M.C.T. was funded by a BBSRC studentship.

REFERENCES

- Bandi, C., Dunn, A. M., Hurst, G. D. D. and Rigaud, T.** (2001). Inherited microorganisms, sex-specific virulence and reproductive parasitism. *Trends in Parasitology* **17**, 88–94.
- Dyer, K. A., Minhas, M. S. and Jaenike, J.** (2005). Expression and modulation of embryonic male-killing in *Drosophila innubila*: opportunities for multilevel selection. *Evolution* **59**, 838–848.
- Gasparich, G. E., Whitcomb, R. F., Dodge, D., French, F. E., Glass, J. and Williamson, D. L.** (2004). The genus *Spiroplasma* and its non-helical descendants: phylogenetic classification, correlation with phenotype and roots of the *Mycoplasma mycoides* clade. *International Journal of Systematic and Evolutionary Microbiology* **54**, 893–918.
- Hasegawa, M., Kishino, H. and Yano, T.** (1985). Dating of the human-ape splitting by a molecular clock of mitochondrial DNA. *Journal of Molecular Evolution* **21**, 160–174.
- Hoffmann, A. A., Turelli, M. and Harshman, L. G.** (1990). Factors affecting the distribution of cytoplasmic incompatibility in *Drosophila simulans*. *Genetics* **126**, 933–948.
- Hunter, M. S., Perlman, S. J. and Kelly, S. E.** (2003). A bacterial symbiont in the *Bacteroidetes* induces cytoplasmic incompatibility in the parasitoid wasp *Encarsia pergandiella*. *Proceedings of the Royal Society of London, B* **270**, 2185–2190.
- Hurst, G. D. D., Bandi, C., Sacchi, L., Cochrane, A., Bertrand, D., Karaca, I. and Majerus, M. E. N.** (1999b). *Adonia variegata* (Coleoptera: Coccinellidae) bears maternally inherited Flavobacteria that kill males only. *Parasitology* **118**, 125–134.
- Hurst, G. D. D., Hammarton, T. M., Bandi, C., Majerus, T. M. O., Bertrand, D. and Majerus, M. E. N.** (1997). The diversity of inherited parasites of insects: the male killing agent of the ladybird beetle *Coleomegilla maculata* is a member of the Flavobacteria. *Genetical Research* **70**, 1–6.
- Hurst, G. D. D. and Jiggins, F. M.** (2000). Male-killing bacteria in insects: mechanisms, incidence, and implications. *Emerging Infectious Diseases* **6**, 329–336.
- Hurst, G. D. D., Jiggins, F. M. and Majerus, M. E. N.** (2003). Inherited microorganisms that selectively kill male hosts: the hidden players of insect evolution? In *Insect Symbiosis* (ed. Bourtzis, K. and Miller, T. A.). CRC Press, Boca Raton, FL, USA.
- Hurst, G. D. D. and Majerus, M. E. N.** (1993). Why do maternally inherited microorganisms kill males? *Heredity* **71**, 81–95.
- Hurst, G. D. D., Majerus, M. E. N. and Walker, L. E.** (1992). Cytoplasmic male killing elements in *Adalia bipunctata* (Linnaeus) (Coleoptera: Coccinellidae). *Heredity* **69**, 84–91.
- Hurst, G. D. D., Schulenburg, J. H. G., Majerus, T. M. O., Bertrand, D., Zakharov, I. A., Baungaard, J., Volkl, W., Stouthamer, R. and Majerus, M. E. N.** (1999a). Invasion of one insect species, *Adalia bipunctata*, by two different male-killing bacteria. *Insect Molecular Biology* **8**, 133–139.
- Hurst, L. D.** (1991). The incidences and evolution of cytoplasmic male killers. *Proceedings of the Royal Society of London, B* **244**, 91–99.
- Jiggins, F. M., Hurst, G. D. D., Jiggins, C. D., Schulenburg, J. H. G. and Majerus, M. E. N.** (2000). The butterfly *Danaus chrysippus* is host to a male-killing *Spiroplasma* bacterium. *Parasitology* **120**, 439–446.
- Kumar, S., Tamura, K. and Nei, M.** (2004). MEGA3: Integrated software for molecular evolutionary genetics

- analysis and sequence alignment. *Briefings in Bioinformatics* **5**, 150–163.
- Majerus, M. E. N.** (1991). Habitat and host plant preferences of British ladybirds. *Entomologists Monthly Magazine* **127**, 167–175.
- Majerus, M. E. N. and Hurst, G. D. D.** (1997). Ladybirds as a model system for the study of male-killing symbionts. *Entomophaga* **42**, 13–20.
- Majerus, M. E. N., Kearns, P. W. E., Ireland, H. and Forge, H.** (1989). Ladybirds as teaching aids: 1 Collecting and culturing. *Journal of Biological Education* **23**, 85–95.
- Majerus, T. M. O., Majerus, M. E. N., Knowles, B., Wheeler, J., Bertrand, D., Kuznetsov, V. N., Ueno, H. and Hurst, G. D. D.** (1998). Extreme variation in the prevalence of inherited male-killing microorganisms between three populations of *Harmonia axyridis* (Coleoptera: Coccinellidae). *Heredity* **81**, 683–691.
- Majerus, T. M. O., Majerus, M. E. N., Schulenburg, J. H. G. and Hurst, G. D. D.** (1999). Molecular identification of a male-killing agent in the ladybird *Harmonia axyridis* (Pallas) (Coleoptera: Coccinellidae). *Insect Molecular Biology* **8**, 551–555.
- Randall, K., Majerus, M. E. N. and Forge, H.** (1992). Characteristics for sex determination in British ladybirds (Coleoptera: Coccinellidae). *Entomologist* **111**, 109–122.
- Schulenburg, J. H. G., Habig, M., Sloggett, J. J., Webberley, K. M., Bertrand, D., Hurst, G. D. D. and Majerus, M. E. N.** (2001). Incidence of male-killing *Rickettsia* spp. (α -proteobacteria) in the ten-spot ladybird beetle *Adalia decempunctata* L. (Coleoptera: Coccinellidae). *Applied and Environmental Microbiology* **67**, 270–277.
- Schulenburg, J. H. G., Hurst, G. D. D., Tetzlaff, D., Booth, G. E., Zakharov, I. A. and Majerus, M. E. N.** (2002). History of infection with different male-killing bacteria in the two-spot ladybird beetle *Adalia bipunctata* revealed through mitochondrial DNA analysis. *Genetics* **160**, 1075–1086.
- Sloggett, J. J. and Majerus, M. E. N.** (2000). Habitat preferences and diet in the predatory Coccinellidae (Coleoptera): an evolutionary perspective. *Biological Journal of The Linnean Society* **70**, 63–88.
- Stevens, L.** (1989). Environmental factors affecting reproductive incompatibility in flour beetles, genus *Tribolium*. *Journal of Invertebrate Pathology* **53**, 78–84.
- Swofford, D. L.** (1998). *PAUP*. Phylogenetic Analysis Using Parsimony (*and Other Methods). Version 4.* Sinauer Associates. Sunderland, Massachusetts.
- van Kuppeveld, F. J. M., van der Logt, H. T. M., van Zoest, M. J., Quint, W. G. V., Niesters, H. G. M., Galama, J. M. D. and Melchers, W. J. G.** (1992). Genus- and species-specific identification of the mycoplasmas by 16s rDNA amplification. *Applied and Environmental Microbiology* **58**, 2606–2615.
- Veneti, Z., Bentley, J. K., Koana, T., Braig, H. R. and Hurst, G. D. D.** (2005). A functional dosage compensation complex required for male killing in *Drosophila*. *Science* **307**, 1461–1463.
- Weeks, A. R., Marec, F. and Breeuwer, J. A. J.** (2001). A mite species that consists entirely of haploid females. *Science* **292**, 2479–2482.
- Werren, J. H. and Windsor, D. M.** (2000). *Wolbachia* infection frequencies in insects: evidence of a global equilibrium? *Proceedings of the Royal Society of London, B* **267**, 1277–1285.
- Werren, J. H., Windsor, D. and Guo, L. R.** (1995). Distribution of *Wolbachia* among neotropical arthropods. *Proceedings of the Royal Society of London, B* **262**, 197–204.
- Zchori-Fein, E. and Perlman, S. J.** (2004). Distribution of the bacterial symbiont *Cardinium* in arthropods. *Molecular Ecology* **13**, 2009–2016.
- Zchori-Fein, E., Gottlieb, Y., Kelly, S. E., Brown, J. K., Wilson, J. M., Karr, T. L. and Hunter, M. S.** (2001). A newly discovered bacterium associated with parthenogenesis and a change in host selection behavior in parasitoid wasps. *Proceedings of the National Academy of Sciences, USA* **98**, 12555–12560.
- Zhou, W. F., Rousset, F. and O'Neill, S.** (1998). Phylogeny and PCR based classification of *Wolbachia* strains using *wsp* gene sequences. *Proceedings of the Royal Society of London, B* **265**, 509–515.